WG15 WINNING GROWTH

MEDICATED

FOR IMPROVED FEED EFFICIENCY IN CATTLE BEING FED IN CONFINEMENT FOR SLAUGHTER

ACTIVE DRUG INGREDIENT

INGREDIENTS

Grain Products, Processed Grain Byproducts, Plant Protein Products, Roughage Products, Molasses, Calcium Carbonate, Salt, Soybean Oil, Potassium Sulfate, Magnesium Sulfate, Magnesium Oxide, Yucca Schidigera Extract, Betaine (Hydrochloride), Saccharomyces Cerevisiae Yeast, Cobalt Carbonate, Lactic Acid, Feugreek Extract, Potassium Chloride, Vitamin E Supplement, Manganous Oxide, Manganese Sulfate, Copper Sulfate, Zinc Oxide, Vitamin A Supplement, Vitamin D3 Supplement, Zinc Sulfate, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, DL Tocopherol-Acetate, Sodium Selenite, Cholecaciferol (D-activated Animal Sterol), Ethylendiamine Dihydriodide, Ethoxyquin (A Preservative), Natural Flavor, Propionic Acid (A Preservative), Sodium Calcium Aluminosilocate, Silicon Dioxide, Mineral Oil.

FEEDING DIRECTIONS & PRECAUTION STATEMENTS; See Reverse

MANUFACTURED BY
MEISS FEED AND SUPPLY
245 S. STATE ROUTE 47, SIBLEY, IL 61773
217-745-2302

NET WEIGHT: 50 LB. BAG

FEEDING DIRECTIONS

This feed is designed as a grower-developer creep feed for show cattle up to 900 lbs, but may be fed to cattle of any age or weight with proper management. Feed complete ration to show cattle at a rate of 4 - 22 lbs per head per day to cattle weighing 150 - 900 lb. to provide 79.0 - 434.5 milligrams of Monensin per head per day It is important to provide available forage as an addition to their daily intake. Provide cattle continuous access to clean, cool water. Store feed in a cool, dry location free of insects, rodents, and other pests. Do not feed if mold or infestation occurs.

CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by horses has been fatal. *Monensin* medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding or mixing errors resulting in high concentrations of *Monensin* could be fatal to cattle and goats. DO NOT exceed the levels of *Monensin* recommended in the feeding directions as reduced average gains may result. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. DO NOT use in calves to be processed for veal.

 $\textbf{CAUTION:} \quad \textbf{Contains copper } \underline{\textbf{DO NOT}} \text{ feed to sheep!}$