

2215RM DAIRY CALF STARTER

MEDICATED

THIS PRODUCT IS FORMULATED AS A STARTER FEED FOR DAIRY CALVES 3 DAYS - 3 MONTHS IN CONFINEMENT. FOR THE PREVENTION AND CONTROL OF COCCIDIOSIS IN CALVES CAUSED BY E. BOVIS AND E. ZUERNII.

ACTIVE DRUG INGREDIENT

MONENSIN.....50.0 g/ton

GUARANTEED ANALYSIS

Crude Protein	Min. 22.0%
Crude Fat	Min. 3.5%
Crude Fiber	Max. 7.5%
Acid Detergent Fiber ADF	Max. 9.0%
Calcium	Min. 0.8% Max. 1.20%
Phosphorus	Min. 0.45%
Selenium	Min. 0.30 ppm
Vitamin A	Min. 9500 IU/LB

INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain Byproducts, Calcium Carbonate, Monocalcium Phosphate, Molasses, Soybean Oil, Corn Starch, Sodium Chloride, Magnesium Oxide, Copper Sulfate, Calcium Sulfate, Manganese Sulfate, Vitamin E Supplement, D-Activated Animal Sterol (A Source of Vitamin D-3), Vitamin A Supplement, Ethylenediamine Dihydriodide, Zinc Sulfate, Potassium Chloride, Potassium Sulfate, Magnesium Sulfate, Cobalt Carbonate, Sodium Selenite, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Mineral Oil, Propionic Acid (A Preservative), Brewers Dried Yeast, Yucca Schidigera Extract, Betaine (Hydrochloride), Lactic Acid, Fenugreek Flavor Extract, Magnesium-Mica, Dried Whey, Dried Bactilus subtilis Fermentation Product, Dried Belfidobacterium Longum Fermentation Product, Dried Bifidobacterium Thermophilum Fermentation Product, Dried Enterococcus Faecium Fermentation Product, Dried Lactobacillus Acidophilus Fermentation Product, Dried Seaweed Meal (Fucaeeae, Banglicaeae, Ulvaceae), Psykkium Seed Husk, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, Natural Flavor, Silcon Dioxide.

FEEDING DIRECTIONS & PRECAUTION STATEMENTS; SEE REVERSE

MANUFACTURED BY

MEISS FEED AND SUPPLY

245 S. STATE ROUTE 47, SIBLEY, IL 61773
217-745-2302

NET WEIGHT: 50 LB. BAG

FEEDING DIRECTIONS

Hand feed continuously to calves (excluding veal calves) at a rate of 1.51 lbs to calves weighing 100 lbs to provide 0.38 mg of monensin per lb of body weight; depending on the severity of the challenge, up to a maximum of 200 mg monensin per head per day. Provide calves continuous access to cool, clean water. Store feed in a cool, dry location free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by horses has been fatal. *Monensin* medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding or mixing errors resulting in high concentrations of *Monensin* could be fatal to cattle and goats. **DO NOT** exceed the levels of *Monensin* recommended in the feeding directions as reduced average gains may result. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. **DO NOT** use in calves to be processed for veal.

CAUTION: Contains copper DO NOT feed to sheep!