

# 1815RM DAIRY CALF GROWER

## MEDICATED

THIS PRODUCT IS FORMULATED AS A FEED FOR DAIRY CALVES 2 - 6 MONTHS OLD. FOR THE PREVENTION AND CONTROL OF COCCIDIOSIS IN CALVES CAUSED BY E. BOVIS AND E. ZUERNI.

### ACTIVE DRUG INGREDIENT

*MONENSIN*.....60.0 g/ton

### GUARANTEED ANALYSIS

Crude Protein ..... Min. 18.0%  
Crude Fat ..... Min. 3.5%  
Crude Fiber ..... Max. 8.0%  
Acid Detergent Fiber (ADF) ..... Max. 12.0%  
Calcium ..... Min. 0.9% Max. 1.40%  
Phosphorus ..... Min. 0.60%  
Selenium ..... Min. 0.60 ppm  
Vitamin A ..... Min. 7500 IU/LB

### INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain Byproducts, Calcium Carbonate, Monocalcium Phosphate, Molasses, Soybean Oil, Corn Starch, Sodium Chloride, Magnesium Oxide, Copper Sulfate, Calcium Sulfate, Manganese Sulfate, Vitamin E Supplement, D-Activated Animal Sterol (A Source of Vitamin D-3), Vitamin A Supplement, Ethylenediamine Dihydriodide, Zinc Sulfate, Potassium Chloride, Potassium Sulfate, Magnesium Sulfate, Cobalt Carbonate, Sodium Selenite, Mineral Oil, Propionic Acid (A Preservative),Brewers Died Yeast, Yucca Schidigera Extract, Betaine (Hydrochloride), Lactic Acid, Fenugreek Flavor Extract, Magnesium-Mica, Dried Whey, Dried Bactilus subtilis Fermentation Product, Dried Beifidobacterium Longum Fermentation Product, Dried Bifidobacterium Thermophilum Fermentation Product, Dried Enterococcus Faecium Fermentation Product,Dried Lactobacillus Acidophilus Fermentation Product, Dried Seaweed Meal (Fucaceae, Banglicaeae, Ulvaceae), Psykkium Seed Husk, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, Natural Flavor, Silcon Dioxide.

**FEEDING DIRECTIONS & PRECAUTION STATEMENTS - See Reverse**

*MANUFACTURED BY*

**MEISS FEED AND SUPPLY**

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

**NET WEIGHT: 50 LB. BAG**

### FEEDING DIRECTIONS

Feed continuously to calves (excluding veal calves) at a rate of 1.65 lbs per 200 lbs of body weight to supply 0.25 mg of Monensin per lb of body weight, depending on the severity of the challenge up to a maximum of 200 mg per head per day. Provide calves continuous access to cool, clean water. Store feed in a cool, dry location free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

**CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT** allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by horses has been fatal. *Monensin* medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding or mixing errors resulting in high concentrations of *Monensin* could be fatal to cattle and goats. **DO NOT** exceed the levels of *Monensin* recommended in the feeding directions as reduced average gains may result. If feed refusals containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. **DO NOT** use in calves to be processed for veal.

**CAUTION:** Contains copper DO NOT feed to sheep!