

1018 LITTLE MOO MIX

MEDICATED

THIS PRODUCT IS FORMULATED AS A STARTER FEED FOR DAIRY & BEEF CALVES. AS AN AID IN THE PREVENTION AND CONTROL OF COCCIDIOSIS IN CALVES CAUSED BY *E. BOVIS* AND *E. ZURNII*.

ACTIVE DRUG INGREDIENT

MONENSIN.....40.0 g/ton

GUARANTEED ANALYSIS

Crude Protein Min. 18.0%
Crude Fat Min. 3.5%
Crude Fiber Max. 8.5%
Calcium Min. 0.8% Max. 1.4%
Phosphorus Min. 0.3%
Selenium Min. 0.308 ppm
Vitamin A Min. 10,500 IU/LB

INGREDIENTS

Grain Products, Plant Protein Products, Roughage Products, Processed Grain Byproducts, Molasses, Monocalcium Phosphate, Calcium Carbonate, Sodium Chloride (Salt), Soybean Oil Magnesium Oxide, Corn Starch, Potassium Sulfate, Magnesium Sulfate, Potassium Chloride, Calcium Sulfate, Manganese Sulfate, Copper Sulfate, Ferrous Carbonate, Zinc Sulfate, Sodium Selenite, Vitamin A Acetate, D-Activated Animal Sterol (A Source of Vitamin D-3), Vitamin E Supplement, Zinc Oxide, Iron Sulfate, Ethylenediamine Dihydriodide, Brewers Dried Yeast, Magnesium-Mica, Yucca Schidigera Extract, Betaine (hydrochloride), Lactic Acid, Fenugreek Flavor Extract, Rice Hulls, Dried Whey, Cobalt Carbonate, Dried Bacillus Subtilis Fermentation Product, Dried Aspergillus Oryzae Fermentation Product, Dried Bifidobacterium Longum Fermentation Product, Dried Bifidobacterium Thermophilum Fermentation Product, Dried Enterococcus Faecium Fermentation Product, Dried Lactobacillus Acidophilus Fermentation Product, Dried Seaweed Meal (Fucaceae, Bangiaceae, Ulvaceae), Psyllium Seed Husk, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, Natural and Artificial Flavors, Ethoxyquin (A Preservative), Propionic Acid, Mineral Oil.

FEEDING DIRECTIONS & PRECAUTION STATEMENTS; SEE REVERSE

MANUFACTURED BY

MEISS FEED AND SUPPLY

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

NET WEIGHT: 50 LB. BAG

FEEDING DIRECTIONS

Provide small amounts of fresh 1018 Little Moo Mix from 4 days after birth. Feed at the rate of 1 lbs per 100 lbs of body weight per day to provide 20 mg of *Monensin* per 100 lbs of body weight. Do not to exceed 100 mg or 5 lbs/100 lbs of body weight. Calf may be weaned off milk when consuming 1.5 lbs of 1018 Little Moo Mix daily, usually 4 - 6 weeks of age. Provide Calves continuous access to cool, clean water. Store feed in a cool, dry location free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by horses has been fatal. *Monensin* medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding or mixing errors resulting in high concentrations of *Monensin* could be fatal to cattle and goats. **DO NOT** exceed the levels of *Monensin* recommended in the feeding directions as reduced average gains may result. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. **DO NOT** use in calves to be processed for veal.

CAUTION: Contains copper DO NOT feed to sheep!