

# 1015 HI-FI CATTLE BUFFET

## MEDICATED

FOR IMPROVED FEED EFFICIENCY IN CATTLE BEING FED IN CONFINEMENT  
FOR SLAUGHTER

---

### ACTIVE DRUG INGREDIENT

*MONENSIN*.....34.0 g/ton

### GUARANTEED ANALYSIS

Crude Protein ..... Min. 15.0%  
Crude Fat ..... Min. 4.0%  
Crude Fiber ..... Max. 21.0%  
Calcium ..... Min. 0.9% Max. 1.3%  
Phosphorus ..... Min. 0.35%  
Selenium ..... Min. 0.18 ppm

### INGREDIENTS

Roughage Products, Grain Products, Processed Grain Byproducts, Plant Protein Products, Corn Starch, Molasses, Calcium Carbonate, Soybean Oil, Salt, Potassium Sulfate, Magnesium Sulfate, Magnesium-Mica, Magnesium Oxide, Yucca Schidigera Extract, Betaine (Hydrochloride), Cobalt Carbonate, Lactic Acid, Fenugreek Extract, Potassium Chloride, Vitamin E Supplement, Manganous Oxide, Manganese Sulfate, Copper Sulfate, Zinc Oxide, Vitamin A Supplement, Vitamin D3 Supplement, Zinc Sulfate, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, DL Tocopherol-Acetate, Sodium Selenite, Cholecalciferol (D-activated Animal Sterol), Ethylendiamine Dihydriodide, Ethoxyquin (A Preservative), Natural Flavor, Propionic Acid, Sodium Calcium Aluminosilicate, Silicon Dioxide, Mineral Oil.

FEEDING DIRECTIONS & PRECAUTION STATEMENTS; SEE REVERSE

MANUFACTURED BY

**MEISS FEED AND SUPPLY**

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

NET WEIGHT: 50 LB. BAG OR BULK INVOICE

### FEEDING DIRECTIONS

Feed complete ration at a rate of 3 - 27 lbs per head per day to cattle weighing 150 - 1300 lb. to provide 51.0 - 459 milligrams of *Monensin* per head per day. Provide cattle continuous access to clean, cool water. Store feed in a cool, dry location free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

**CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT** allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by horses has been fatal. *Monensin* medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding or mixing errors resulting in high concentrations of *Monensin* could be fatal to cattle and goats. **DO NOT** exceed the levels of *Monensin* recommended in the feeding directions as reduced average gains may result. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. **DO NOT** use in calves to be processed for veal.

**CAUTION:** Contains copper DO NOT feed to sheep!