

# 1013 CATTLE FINISHER

## MEDICATED

FOR IMPROVED FEED EFFICIENCY IN CATTLE BEING FED IN CONFINEMENT FOR SLAUGHTER

### ACTIVE DRUG INGREDIENT

MONENSIN.....31.0 g/ton

### GUARANTEED ANALYSIS

Crude Protein ..... Min. 12.0%  
Crude Fat ..... Min. 3.0%  
Crude Fiber ..... Max. 15.0%  
Selenium ..... Min. 0.30 ppm

### INGREDIENTS

Grain Products, Processed Grain Byproducts, Roughage Products, Calcium Carbonate, Molasses, Soybean Oil, Salt, Manganese Sulfate, Zinc Sulfate, Magnesium Oxide, Ferrous Sulfate, Copper Sulfate, Zinc Oxide, DL-Alpha Tocopheryl Acetate, Potassium Chloride, Sodium Selenite, Magnesium Sulfate, Potassium Sulfate, Thiamine Mononitrate, Ethylenediamine Dihydriodide, Vitamin A Acetate, Cholecalciferol (D-Activated Animal Sterol), Cobalt Carbonate, Magnesium-Mica, Yucca Schidigera Extract, Betaine (Hydrochloride), Lactic Acid, Fenugreek Extract, Vitamin E Supplement, Manganous Oxide, Vitamin A Supplement, Vitamin D3 Supplement, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, Natural Flavor, Propionic Acid, Sodium Calcium Aluminosilicate, Silicon Dioxide, Ethoxyquin (A Preservative), Mineral Oil.

### FEEDING DIRECTIONS & PRECAUTION STATEMENTS; SEE REVERSE

### MANUFACTURED BY

### MEISS FEED AND SUPPLY

245 S. STATE ROUTE 47, SIBLEY, IL 61773  
217-745-2302

**NET WEIGHT: 50 LB. BAG OR BULK INVOICE**

#### FEEDING DIRECTIONS

Feed complete ration at a rate of 16 -29 lbs per head per day to cattle weighing 800 - 1350 lbs. to provide 248.0 - 449.5 milligrams of *Monensin* per head per day. It is important to provide hay as an addition to their daily intake. Store feed in a cool, dry location free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

**CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT** allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by horses has been fatal. *Monensin* medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of *Monensin* recommended in the feeding directions, as reduced average gains may result. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. **DO NOT** use in calves to be processed for veal.

CAUTION: Contains copper DO NOT feed to sheep!