

1011 FINAL FINISHER

MEDICATED

FOR IMPROVED FEED EFFICIENCY IN CATTLE BEING FED IN
CONFINEMENT FOR SLAUGHTER

ACTIVE DRUG INGREDIENT

MONENSIN 30.0 g/ton

GUARANTEED ANALYSIS

Crude Protein Min. 11.0%

(This includes not more than 1.6% equivalent crude protein from non-protein nitrogen)

Crude Fat Min. 3.0%

Crude Fiber Max. 4.0%

Selenium Min. 0.15 ppm

INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain Byproducts, Calcium Carbonate, Urea, Potassium Chloride, Monocalcium Phosphate, Yeast Culture, Calcium Sulfate, Magnesium Sulfate, Potassium Sulfate, Salt, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Ferrous Sulfate, DL-Alpha Tocopheryl Acetate, Sodium Selenite, Mineral Oil, Vitamin A Supplement, D-Activated Animal Sterol (Source of Vitamin D-3), Ethylenediamine Dihydriodide, Ethoxyquin (A Preservative), Cobalt Carbonate, Animal & Vegetable Fat, Sodium Calcium Aluminosilicate, Natural Flavor.

FEEDING DIRECTIONS & PRECAUTION STATEMENTS; SEE REVERSE

MANUFACTURED BY

MEISS FEED AND SUPPLY

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

NET WEIGHT: 50 LB. BAG OR BULK INVOICE

FEEDING DIRECTIONS

Feed complete ration at a rate of 8 - 28 lbs per head per day to cattle weighing 400 - 1450 lbs. to provide 120 - 420 milligrams of *Monensin* per head per day. It is important to provide hay as an addition to their daily intake. Store feed in a cool, dry location free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by horses has been fatal. *Monensin* medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of *Monensin* recommended in the feeding directions, as reduced average gains may result. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. **DO NOT** use in calves to be processed for veal.

CAUTION: Contains copper DO NOT feed to sheep!