

4080 BEEF SUPPLEMENT

MEDICATED

FOR IMPROVED FEED EFFICIENCY IN CATTLE BEING FED IN CONFINEMENT FOR SLAUGHTER. FOR THE PREVENTION AND CONTROL OF COCCIDIOSIS CAUSED BY EIMERIA BOVIS AND E. ZUERHII IN CATTLE MAINTAINED IN CONFINEMENT

ACTIVE DRUG INGREDIENT

MONENSIN..... 300 g/ton

GUARANTEED ANALYSIS

Crude Protein Min. 40.0%
(This includes not more than 15.0% equivalent crude protein from non-protein nitrogen)
Crude Fat Min. 1.0%
Crude Fiber Max. 10.0%
Calcium Min. 3.5% Max. 4.5%
Phosphorus Min. 0.8%
Vitamin A Min. 22,000 IU/LB

INGREDIENTS

Plant Protein Products, Processed Grain Byproducts, Roughage Products, Calcium Carbonate, Urea, Potassium Chloride, Monocalcium Phosphate, Yeast Culture, Calcium Sulfate, Magnesium Sulfate, Potassium Sulfate, Salt, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Ferrous Sulfate, DL-Alpha Tocopheryl Acetate, Sodium Selenite, Mineral Oil, Vitamin A Supplement, D-Activated Animal Sterol (Source of Vitamin D-3), Ethylenediamine Dihydriodide, Cobalt Carbonate, Animal & Vegetable Fat, Sodium Calcium Aluminosilicate, Ethoxyquin (A Preservative).

MIXING/FEEDING DIRECTIONS & PRECAUTION STATEMENTS; *SEE REVERSE*

MANUFACTURED BY

MEISS FEED AND SUPPLY

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

NET WEIGHT: 50 LB.BAG OR BULK INVOICE

MIXING DIRECTIONS

Thoroughly mix 4080 Beef Supplement with grain or complete feed before feeding. **DO NOT FEED ALONE.** Mix 150 lbs. of 4080 Beef Supplement with 1850 lbs. of feed or feed ingredients to make one ton of *Type C Medicated Feed* containing 22.5 grams of *Monensin* per ton.

FEEDING DIRECTIONS

Feed the complete ration at the rate of 10 - 29 lbs./head/day to cattle weighing 800 lbs, to provide 112.5 - 326.25 milligrams of *Monensin* per head per day. Provide cattle continuous access to clean, cool water. Store feed in a cool, dry location free of insects, rodents, and other pests. **Do not feed if mold or infestation occurs.**

CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by equines has been fatal. *Monensin* medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. **DO NOT FEED UNDILUTED.** Feeding undiluted, or mixing errors resulting in high concentrations of *Monensin* has been *fatal* to cattle. **DO NOT** exceed the levels of *Monensin* recommended in the feeding directions, as reduced average daily gains may result. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. Warning a withdrawal time has not been established for pre-ruminating calves. **DO NOT** use in calves to be processed for veal.

CAUTION: Contains copper **DO NOT** feed to sheep!

Rev: 2-3-17