

1084 BEEF BALANCER

MEDICATED

FOR IMPROVED FEED EFFICIENCY

ACTIVE DRUG INGREDIENT

MONENSIN..... 1000 g/ton

GUARANTEED ANALYSIS

Calcium Min. 30.0% Max. 32.0%
Salt Min. 14.1% Max. 15.6%
Selenium Min. 13.0 ppm
Vitamin A Min. 103,000 IU/LB
Vitamin D-3 Min. 30,000 IU/LB
Vitamin E Min. 403.5 IU/LB

INGREDIENTS

Calcium Carbonate, Salt, Manganese Sulfate, Zinc Sulfate, Soybean Oil, Magnesium Oxide, Ferrous Sulfate, Copper Sulfate, Zinc Oxide, DL-Alpha Tocopheryl Acetate, Potassium Chloride, Sodium Selenite, Magnesium Sulfate, Potassium Sulfate, Thiamine Mononitrate, Ethylenediamine Dihydriodide, Vitamin A Acetate, Cholecalciferol (D-Activated Animal Sterol), Cobalt Carbonate.

MIXING/FEEDING DIRECTIONS & PRECAUTION STATEMENTS; SEE REVERSE

MANUFACTURED BY

MEISS FEED AND SUPPLY

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

NET WEIGHT: 50 LB. BAG OR BULK INVOICE

MIXING DIRECTIONS

Thoroughly mix Beef Corn Byproduct Balancer 1084 with grain or complete feed before feeding. **DO NOT FEED ALONE.** Mix 50 lbs. of Beef Corn Byproduct Balancer 1084 with 1950 lbs. of feed or feed ingredients to make one ton of *Type C Medicated Feed* containing 25 grams of *Monensin* per ton.

FEEDING DIRECTIONS

Feed the complete ration at the rate of 4 - 28 lb. per head per day to provide 50 - 350 milligrams of *Monensin* per head per day. Provide cattle continuous access to clean, cool water. Store feed in a cool, dry location free of insects, rodents, and other pests. ***Do not feed if mold or infestation occurs.***

CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT allow horses or other equines access to formulations containing *Monensin*. Ingestion of *Monensin* by equines has been fatal. *Monensin* medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. DO NOT exceed the levels of *Monensin* recommended in the feeding directions, as reduced average daily gains may result. Feeding, or mixing errors resulting in high concentrations of *Monensin* could be *fatal* to cattle: must be thoroughly mixed in feeds before use: DO NOT FEED UNDILUTED. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. DO NOT use in calves to be processed for veal.

CAUTION: Follow label instructions! The additions to feed at higher levels of this product containing selenium is not permitted!

CAUTION: Contains copper DO NOT feed to sheep!