

3218 KID STARTING BUFFET

MEDICATED

FOR PREVENTION OF COCCIDIOUS CAUSED BY EIMERIA CRANDALLIS,
E. CHRISTENSENI AND E. NINAKOHLIYAKIMOVAE

ACTIVE DRUG INGREDIENT

MONENSIN.....20.0 g/ton

GUARANTEED ANALYSIS

Crude Protein Min. 18.0%
(This includes not more than 1.5% equivalent crude protein from non-protein nitrogen)
Crude Fat Min. 5.0%
Crude Fiber Max. 9.0%
Calcium Min. 0.8% Max. 1.8%
Phosphorus Min. 0.45%
Salt Min. 0.8% Max. 1.2%
Sodium Min. 0.31% Max. 0.52%
Selenium Min. 0.18 ppm

INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain Byproducts, Molasses Products, Roughage Products, Soybean Oil, Monocalcium Phosphate, Calcium Carbonate, Sodium Chloride (Salt), Ammonium Chloride, Magnesium Oxide, Corn Starch, Potassium Sulfate, Magnesium Sulfate, Potassium Chloride, Calcium Sulfate, Manganese Sulfate, Copper Sulfate, Ferrous Carbonate, Zinc Sulfate, Sodium Selenite, Vitamin A Acetate, D-Activated Animal Sterol (A Source of Vitamin D-3), Vitamin E Supplement, Zinc Oxide, Iron Sulfate, Ethylenediamine Dihydrochloride, Brewers Dried Yeast, Magnesium-Mica, Yucca Schidigera Extract, Betaine (hydrochloride), Lactic Acid, Fenugreek Flavor Extract, Thiamine Mononitrate, Rice Hulls, Dried Whey, Cobalt Carbonate, Dried Bacillus Subtilis Fermentation Product, Dried Aspergillus Oryzae Fermentation Product, Dried Bifidobacterium Longum Fermentation Product, Dried Bifidobacterium Thermophilum Fermentation Product, Dried Enterococcus Faecium Fermentation Product, Dried Lactobacillus Acidophilus Fermentation Product, Dried Seaweed Meal (Fucaceae, Bangiaceae, Ulvaceae), Psyllium Seed Husk, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, Natural and Artificial Flavors, Propionic Acid, Mineral Oil.

FEEDING DIRECTIONS & PRECAUTION STATEMENTS: SEE REVERSE

MANUFACTURED BY

MEISS FEED AND SUPPLY

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

NET WEIGHT: 50 LB BAG OR BULK INVOICE

FEEDING DIRECTIONS

Feed continuously to newly weaned, growing kids in confinement. It is important to provide hay as an addition to their daily intake. Provide kids continuous access to clean, cool water. Store feed in a cool, dry location free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN: DO NOT allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by horses has been fatal. *Monensin* medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding or mixing errors resulting in high concentrations of *Monensin* could be fatal to cattle and goats. DO NOT exceed the levels of *Monensin* recommended in the feeding directions as reduced average gains may result. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. DO NOT use in calves to be processed for veal. Do not feed to lactating goats.

CAUTION: Contains copper DO NOT feed to sheep!