

3214 FINAL BUFFET

MEDICATED

FOR PREVENTION OF COCCIDIOUSIS CAUSED BY EIMERIA CRANDALLIS, E.
CHRISTENSENI AND E. NINAKOHLIYAKIMOVAE

ACTIVE DRUG INGREDIENT

MONENSIN.....20 g/ton

GUARANTEED ANALYSIS

Crude Protein Min. 14.0%
(This includes not more than 1.5% equivalent crude protein from non-protein nitrogen)
Crude Fat Min. 4.5%
Crude Fiber Max. 16.0%
Calcium Min. 0.8% Max. 1.8%
Phosphorus Min. 0.45%
Salt Min. 0.8% Max. 1.2%
Sodium Min. 0.31% Max. 0.52%
Selenium Min. 0.20 ppm

INGREDIENTS

Grain Products, Processed Grain Byproducts, Plant Protein Products, Roughage Products, Molasses Products, Soybean Oil, Corn Starch, Calcium Carbonate, Salt, Ammonium Chloride, Potassium Sulfate, Magnesium Sulfate, Magnesium-Mica, Magnesium Oxide, Yucca Schidigera Extract, Betaine (Hydrochloride), Copper Sulfate, Cobalt Carbonate, Lactic Acid, Fenugreek Extract, Potassium Chloride, Vitamin E Supplement, Vitamin D-3 Supplement, Zinc Sulfate, Thiamine Mononitrate, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, DL Tocopherol-Acetate, Sodium Selenite, Cholecalciferol (D-activated Animal Sterol), Ethylendiamine Dihydriodide, Natural Flavor, Propionic Acid, Sodium Calcium Aluminosilocate, Silicon Dioxide, Mineral Oil.

FEEDING DIRECTIONS & PRECAUTION STATEMENTS: SEE REVERSE

MANUFACTURED BY

MEISS FEED AND SUPPLY

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

NET WEIGHT: 50 LB BAG OR BULK INVOICE

FEEDING DIRECTIONS

Feed continuously to growing kids fed in confinement once goats average 60% of intended market weight. It is important to provide hay as an addition to their daily intake. Provide kids continuous access to clean, cool water. Store feed in a cool, dry location free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

CAUTION FOR ALL PRODUCTS CONTAINING MONENSIN:

DO NOT allow horses or other equines access to feeds containing *Monensin*. Ingestion of *Monensin* by horses has been fatal. *Monensin* medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding or mixing errors resulting in high concentrations of *Monensin* could be fatal to cattle and goats. DO NOT exceed the levels of *Monensin* recommended in the feeding directions as reduced average gains may result. If feed refusals fed containing *Monensin* are fed to other groups of cattle, the concentration of *Monensin* in the refusals and the amount of the refusals fed should be taken into consideration to prevent *Monensin* overdosing. A withdrawal time has not been established for pre-ruminating calves. DO NOT use in calves to be processed for veal. Do not feed to lactating goats.

CAUTION: Contains copper DO NOT feed to sheep!