

2818 STARTING LAMB BUFFET MEDICATED

FOR PREVENTION OF COCCIDIOSIS CAUSED BY *E. OVINA*, *E. CRANDALLIS*, *E. OVINOIDALIS* (*E. NINAKOHLIYAKIMOVAE*), *E. PARVA*, AND *E. ITRICATA* IN SHEEP MAINTAINED IN CONFINEMENT

ACTIVE DRUG INGREDIENT

Lasalosisid 30.0 g/ton

GUARANTEED ANALYSIS

Crude Protein Min. 18.0%

(This includes not more than 1.7% equivalent crude protein from Non-Protein Nitrogen)

Crude Fat Min. 4.0%

Crude Fiber Max.

18.0%

Calcium Min. 1.0% Max. 1.5%

Phosphorus Min. 0.35%

Salt Min. 0.8% Max. 1.3%

Sodium Min. 0.31% Max. 0.82%

Selenium Min. 0.30 ppm

INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain Bypoducts, Roughage Products, Corn Starch, Calcium Carbonate, Salt, Ammonium Chloride, Potassium Sulfate, Magnesium Sulfate, Magnesium-Mica, Magnesium Oxide, Molybdenum, Yucca Schidigera Extract, Betaine (Hydrochloride), Cobalt Carbonate, Lactic Acid, Fenugreek Extract, Potassium Chloride, Vitamin E Supplement, Vitamin D-3 Supplement, Zinc Sulfate, Thiamine Mononitrate, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, DL Tocopherol-Acetate, Sodium Selenite, Cholecalciferol (D-activated Animal Sterol), Ethylendiamine Dihydriodide, Ethoxyquin (A Preservative), Natural Flavor, Molasses, Soybean Oil, Propionic Acid, Sodium Calcium Aluminosilicate, Silicon Dioxide, Mineral Oil.

FEEDING DIRECTIONS & PRECAUTION STATEMENTS; SEE REVERSE

MANUFACTURED BY

MEISS FEED AND SUPPLY

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

NET WEIGHT: 50 LB BAG OR BULK INVOICE

FEEDING DIRECTIONS

Feed continuously to lambs prior to weaning, continuing up to 3 weeks after final weaning date at a rate of 1.0 – 4.67 lbs per head per day, to provide not less than 15 mg and not more than 70 mg of lasalocid per head per day depending on body weight. Be sure to provide continuous access to clean, cool water. Store feed in a cool, dry location, free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

Caution: Do not allow horses or other equines access to premixes or supplements containing lasalocid, as ingestion may be fatal. The safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors resulting in excessive concentrations of lasalocid could be fatal to cattle and sheep.

Caution: Follow label directions. The addition to feed of higher levels of this product containing selenium is not permitted. The feeding of this product should not exceed the recommended feeding levels.