

# 2828 LACTATING BUFFET MEDICATED

FOR PREVENTION OF COCCIDIOSIS CAUSED BY *E. OVINA*, *E. CRANDALLIS*, *E. OVINOIDALIS* (*E. NINAKOHLIYAKIMOVAE*), *E. PARVA*, AND *E. ITRICATA* IN SHEEP MAINTAINED IN CONFINEMENT

---

#### ACTIVE DRUG INGREDIENT

Lasalosid ..... 30.0 g/ton

#### GUARANTEED ANALYSIS

Crude Protein ..... Min. 18.0%  
Crude Fat ..... Min. 4.5%  
Crude Fiber ..... Max. 20.0%  
Calcium ..... Min. 1.0% Max. 1.5%  
Phosphorus ..... Min. 0.35%  
Salt ..... Min. 0.9% Max. 1.4%  
Sodium ..... Min. 0.31% Max. 0.81%  
Selenium ..... Min. 0.30 ppm

#### INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain Byproducts, Roughage Products, Molasses, Soybean Oil, Calcium Carbonate, Salt, Corn Starch, Potassium Sulfate, Magnesium Sulfate, Magnesium-Mica, Magnesium Oxide, Molybdenum, Yucca Schidigera Extract, Betaine (Hydrochloride), Cobalt Carbonate, Lactic Acid, Fenugreek Extract, Potassium Chloride, Vitamin E Supplement, Vitamin D-3 Supplement, Zinc Sulfate, Thiamine Mononitrate, Chicory Root, Red Pepper, Anise Oil, Cloves, Saccharin Sodium, Hemicellulose Extract, Origanum Oil, Lemongrass Oil, DL Tocopherol-Acetate, Sodium Selenite, Cholecalciferol (D-activated Animal Sterol), Ethylenediamine Dihydriodide, Ethoxyquin (A Preservative), Natural Flavor, Propionic Acid, Sodium Calcium Aluminosilicate, Silicon Dioxide, Mineral Oil.

FEEDING DIRECTIONS & PRECAUTION STATEMENTS; SEE REVERSE

MANUFACTURED BY

**MEISS FEED AND SUPPLY**

245 S. STATE ROUTE 47, SIBLEY, IL 61773

217-745-2302

NET WEIGHT: 50 LB BAG OR BULK INVOICE

#### FEEDING DIRECTIONS

Feed continuously to ewes up to 3 weeks prior to lambing, and ewes with nursing lambs at a rate of 1.00 - 4.67 lbs per head per day, to provide not less than 15 mg and not more than 70 mg of lasalocid per head per day depending on body weight. Be sure to provide continuous access to clean, cool water. Store feed in a cool, dry location, free of insects, rodents, and other pests. *Do not feed if mold or infestation occurs.*

**Caution:** Do not allow horses or other equines access to premixes or supplements containing lasalocid, as ingestion may be fatal. The safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors resulting in excessive concentrations of lasalocid could be fatal to cattle and sheep.

**Caution:** *Follow label directions. The addition to feed of higher levels of this product containing selenium is not permitted. The feeding of this product should not exceed the recommended feeding levels.*